



Zn²⁺ modulation of ATP-responses at recombinant P2X₂ receptors and its dependence on extracellular pH

S.S. Wildman,¹ B.F. King & G. Burnstock

Department of Anatomy & Developmental Biology, University College London, Gower Street, London WC1E 6BT

1 Using recombinant P2X₂ receptors expressed in *Xenopus* oocytes, the modulatory effects of zinc (Zn²⁺) on ATP-responses were studied under voltage-clamp conditions and at different levels of extracellular pH.

2 Zn²⁺ (0.3–300 μM) added to the bathing medium potentiated ATP-activated membrane currents, increasing ATP-responses by up to 20 fold. This potentiating effect was reversed on washout. Zn²⁺-potentiation was reduced in an exponential manner (decaying 1/e in 42 s) as the interval was lengthened between adding Zn²⁺ then ATP to the superfusate.

3 The potentiating effect of Zn²⁺ was progressively diminished by acidic shifts in extracellular pH (pH_e) which, of itself, also potentiated ATP-responses at P2X₂ receptors. The maximal potentiating effects of Zn²⁺ and H⁺ were not additive.

4 Neither Zn²⁺ nor H⁺ potentiation of ATP-responses was abolished by diethylpyrocarbonate (DEPC, 0.3–3 mM), which irreversibly denatures histidyl residues. Nine histidyl residues are present in the extracellular loop of P2X₂ receptors.

5 Zn²⁺ also enhanced the blocking activity of the P2 receptor antagonist suramin at P2X₂ receptors. Therefore, Zn²⁺ also mimics H⁺ in increasing suramin-activity at P2X₂ receptors.

6 In summary, Zn²⁺ and H⁺ potentiate agonist and antagonist activity at P2X₂ receptors but their effects are not wholly alike for receptor agonism. There, the potentiating effects of Zn²⁺ are time-dependent and gradually convert to inhibition while those of H⁺ are time-independent, persistent and more potent, suggesting that either these modulators interact in a different way with a single allosteric site or with different allosteric sites.

Keywords: Zinc; extracellular pH; ATP; P2X receptor; oocyte

Introduction

Adenosine 5'-triphosphate (ATP) acts as a fast neurotransmitter for a family of ligand-gated ion channels, the P2X receptors, of which seven subtypes (P2X₁₋₇) have been cloned from various tissues, including central and peripheral neurones, cardiac and smooth muscle, secretory epithelia and immune cells (for reviews, see: Burnstock, 1996; Buell *et al.*, 1996; Burnstock & King, 1996; North, 1996). P2X₁₋₇ subunits share a common topology of intracellular N- and C-termini, two transmembrane spanning regions and a large extracellular loop (with 10 conserved cysteine residues) which is presumed to form the ATP binding pocket (North, 1996). The stoichiometry of cloned P2X subtypes and native P2X receptors is unknown (Buell *et al.*, 1996).

Only one member of this ATP receptor/ion-channel family, the P2X₂ receptor, is strongly affected by extracellular Zn²⁺ which potentiates ATP-responses at recombinant P2X₂ receptors (Brake *et al.*, 1994; Nakazawa & Ohno, 1996; 1997). Zn²⁺ also potentiates ATP-activated currents and ATP-evoked dopamine release from rat pheochromocytoma PC12 cells (Koizumi *et al.*, 1995) as well as potentiating ATP-activated currents in rat sensory (nodose) and sympathetic (SCG) neurones (Cloues *et al.*, 1993; Li *et al.*, 1993; 1996; Cloues, 1995). Since transcripts for the P2X₂ subunit were isolated from rat PC12 cells (Brake *et al.*, 1994) and are also present in rat sensory neurones (Lewis *et al.*, 1995) and rat sympathetic (SCG) neurones (Collo *et al.*, 1996), a strong

potentiating effect by Zn²⁺ may implicate the inclusion of P2X₂ subunits in these native P2X receptors.

Agonist activity at the P2X₂ receptor is also strongly affected by extracellular pH (pH_e), with acidification increasing and alkalization decreasing agonist potency at this ATP-gated ion channel (King *et al.*, 1996c; 1997; Wildman *et al.*, 1997). A similar modulatory effect of pH_e was also found for ATP-responses at native P2X receptors in rat sensory neurones (Li *et al.*, 1996). In the present study, we have investigated the modulatory effects of Zn²⁺ on ATP activity at the P2X₂ receptor and explored the relationship between pH_e and Zn²⁺ potentiation to see if Zn²⁺ and H⁺ ions either act separately or by a common mechanism at P2X₂ receptors.

Methods

Oocyte preparation

Xenopus laevis were anaesthetized with Tricaine (0.1% w/v) and killed by decapitation. Ovarian lobes were excised and stored (at 4°C) in a Barth's solution (pH 7.45) containing (mM): NaCl 110, KCl 1, NaHCO₃ 2.4, Tris-HCl 7.5, Ca(NO₃)₂ 0.33, CaCl₂ 0.41, MgSO₄ 0.82 and gentamycin sulphate 50 μg l⁻¹. Mature oocytes (stages V and VI) were defolliculated by a two-step process involving collagenase treatment (Type 1A, 2 mg ml⁻¹ in a Ca²⁺-free Ringer, for 2–3 h) then stripping away the follicle cell layer with fine forceps. Defolliculation removes the native P1 and P2 receptors on the follicle cell layer that envelops oocytes (King *et al.*, 1996a; b). Defolliculated oocytes were injected cytosolically with rat

¹ Author for correspondence at: Autonomic Neuroscience Institute, Royal Free Hospital School of Medicine, Rowland Hill Street, Hampstead, London NW3 2PF.

P2X₂ cRNA (40 nl, 0.1–1 µg ml⁻¹), then incubated at 18°C in Barth's solution for 48 h to allow full receptor expression, and stored at 4°C in Barth's solution for up to 12 days until used in electrophysiological experiments.

Electrophysiology

ATP-activated currents were measured from cRNA-injected oocytes held under voltage-clamp with a twin-electrode amplifier (Axoclamp 2B, with HS-2A head-stages). The voltage-recording (1–2 MΩ tip resistance) and current-recording (5 MΩ tip resistance) microelectrodes were filled with 0.6 M K₂SO₄ and 3.0 M KCl, respectively. Oocytes were placed in an electrophysiological chamber (0.5 ml vol) and superfused (5 ml min⁻¹, at 18°C) with a Ringer solution containing (mM): NaCl 110, KCl 2.5, HEPES 5, CaCl₂ 1.8; adjusted to pH 7.5. Where necessary, the pH of the bathing Ringer solution (pH_e) was adjusted with either 1.0 N HCl or 1.0 N NaOH, to achieve the desired acidic or alkaline shift. Electrophysiological data were filtered initially at 3 kHz, captured at a rate of 20 Hz on a computer with the MP100WSW interface (Biopac Systems, Inc.) and analysed by a commercial software package (Acqknowledge III, Biopac).

ATP-containing solutions

ATP was added to the Ringer solution (at the final concentrations given in the text) and superfused over dual-impaled oocytes by a gravity-fed continuous flow system, which allowed the rapid addition and washout of the agonist. ATP was applied for 120 s or until currents reached a plateau, then washed off with normal Ringer solution for a period of 5 min (or for times indicated in the text). The P2X₂ receptor is a slowly-desensitizing receptor and recovers rapidly (within 5 min) from agonist activation (Brake *et al.*, 1994). For ATP concentration-response (C/R) curves, data were normalized to the maximum current (*I*_{max}) evoked by ATP (100 µM) at pH 7.5. The concentration that evoked 50% of the maximum response (EC₅₀) was taken from Hill plots constructed with the transform log (*I*/*I*_{max} - *I*), where *I* is the current evoked by each concentration of agonist. The Hill coefficient (n_H) was taken from the slope of Hill plots. Concentration-response curves for ATP shown in Results were fitted to the Hill equation by use of commercial software (Prism v1.03, GraphPad).

Zinc-containing solutions

The modulatory activity of Zn²⁺ at the P2X₂ receptor was tested in two ways. In most experiments, Zn²⁺ was added to ATP solutions (at the final concentrations given in the text) and both applied simultaneously to oocytes. In other experiments, Zn²⁺ was added to the Ringer solution and oocytes were pre-incubated for the times mentioned in the text before ATP was applied. In some experiments, the transition metal was added to the superfusate and P2X₂ receptors were superfused for 60 min before ATP concentration-response curves were constructed. In such experiments, Zn²⁺ was not removed from the superfusate during agonist washout. Data on Zn²⁺ potentiation of ATP-responses were normalized to control responses to 100 µM ATP (a supramaximal stimulus at pH 7.5). EC₅₀ and n_H values for Zn²⁺ concentration-response curves were calculated from Hill plots, with the maximum effect of Zn²⁺ at each level of pH_e in calculations. Diethylpyrocarbonate (DEPC) was used in some experiments to denature histidyl residues (Miles, 1977) on the extracellular loop of P2X₂ receptors. Oocytes were pre-incubated for 10 min

in Ringer solution containing DEPC (0.3 mM–3 mM) then washed with normal Ringer before ATP activity was retested in the presence of Zn²⁺ and under acidic conditions.

Statistics

Data are presented as mean ± s.e.mean of 4 sets of data from different oocyte batches. Significant differences were determined by unpaired Student's *t* test, by use of commercial software (Instat v2.05A, GraphPad).

Chemicals

Common salts and reagents were AnalaR grade (Aldrich Chemicals, U.K.). ATP disodium salt, diethylpyrocarbonate (DEPC) and zinc chloride were purchased from Sigma Chemical Co. (Dorset, U.K.). Suramin was a gift from Bayer plc (U.K.).

Results

Zn²⁺ potentiation of *I*_{ATP}

When Zn²⁺ (0.3–300 µM) and ATP (3 µM, EC₅₀) were added simultaneously to the superfusate (T = 0 min), ATP-responses at P2X₂ receptors were potentiated 2 fold at 0.3 µM Zn²⁺ and 15 to 20 fold at 30 µM, after which higher concentrations (50–300 µM) failed to enhance ATP activity further (Figure 1a). At




Figure 1 Zn²⁺ potentiation of *I*_{ATP} at P2X₂ receptors. Whole-cell membrane currents in *Xenopus* oocytes expressing P2X₂ receptors and activated by ATP (3 µM, EC₅₀ at pH 7.5) were potentiated in a concentration-dependent and time-dependent manner by Zn²⁺ (0.3–300 µM) added to the superfusate. In (a), Zn²⁺ and ATP were applied at the same time to oocytes while in (b), Zn²⁺ was applied continuously for 5 min before the application of ATP. Holding potential (*V*_h) was –80 mV; data in (a) and (b) from two separate oocytes.

high concentrations (300 μM and greater), the potentiating effect of Zn²⁺ appeared to wane slightly (see Figures 1a, 3a). The EC₅₀ value for Zn²⁺ potentiation was $9.9 \pm 1.9 \mu\text{M}$ ($n=4$) at pH 7.5 and this effect reversed on washout. When Zn²⁺ (0.3–300 μM) was applied 5 min before ATP ($T=5$ min), ATP-responses were unaffected by 1 μM Zn²⁺ and potentiated 10 to 20 fold at 30 μM , after which higher concentrations (50–300 μM) progressively inhibited ATP activity (Figure 1b). In this case, the apparent EC₅₀ value for Zn²⁺ potentiation was $6.1 \pm 1.2 \mu\text{M}$ ($n=4$) at pH 7.5 (not significantly different from $T=0$ data). Both the potentiation and inhibitory effects of Zn²⁺ were reversed on washout. The inhibitory effect was dependent on the length of Zn²⁺ pre-incubation but not Zn²⁺ concentration. For 300 μM Zn²⁺ (where the inhibitory effect was most profound), ATP-responses were initially potentiated when Zn²⁺ was applied simultaneously with ATP (Figure 2a). However, the degree of potentiation was progressively reduced when the pre-incubation period for Zn²⁺ was lengthened from 30 s to 300 s (Figure 2a). Averaged data for this run-down could be fitted to an exponential function, the time constant (decay by $1/e$) for which was 42 ± 3 s ($n=4$) (Figure 2b).




Figure 2 Run-down of Zn²⁺ potentiation of I_{ATP} at P2X₂ receptors. Whole-cell membrane currents to ATP and potentiated by Zn²⁺ (300 μM) showed run-down which was dependent on the duration of pre-incubation with the transition metal before the addition of the agonist. In (a), ATP (5 μM , at pH 7.5) produced a submaximal response ($V_h = -80$ mV) which was potentiated by Zn²⁺ (300 μM) added simultaneously with the agonist ($T=0$). The amplitude of the ATP-response progressively decreased as the pre-incubation time with Zn²⁺ (300 μM) was increased from 30 s to 300 s. In (b), the relationship between the amplitude of I_{ATP} potentiated by Zn²⁺ (300 μM) and the duration of Zn²⁺ pre-incubation. The time constant (τ) for run-down by $1/e$ was 42 ± 3 s ($n=4$). The curve was fitted for exponential decay by use of commercial software (Prism v1.03, GraphPad).

Effect of pH on Zn²⁺ potentiation of I_{ATP}

The potentiating effect of Zn²⁺ on ATP-responses was dependent on extracellular pH, regardless of the length of Zn²⁺ pre-incubation. Where Zn²⁺ was applied simultaneously with ATP, the maximum potentiating effect of Zn²⁺ on ATP-response was reduced by 60% at pH 6.5 and abolished at pH 5.5 (Figure 3a). The potentiating effect of Zn²⁺ was not significantly different at pH 8.0 compared to its effect at pH 7.5 (Figure 3a). The EC₅₀ values for the Zn²⁺ potentiation of I_{ATP} were ($n=4$): pH 8.0, $11.4 \pm 1.7 \mu\text{M}$; pH 7.5, $9.9 \pm 1.9 \mu\text{M}$; pH




Figure 3 pH-dependence of Zn²⁺ potentiation of I_{ATP} . Concentration-response curves for the potentiating effect of Zn²⁺ (0.3–300 μM) on whole-cell membrane currents to ATP were studied at different levels of extracellular pH. In (a), Zn²⁺ and ATP were applied at the same time ($T=0$ min) to oocytes while in (b), Zn²⁺ was applied continuously for 5 min ($T=5$ min) before the application of ATP. Control ATP-responses were evoked by the EC_{5%} (i.e. control ATP-responses were amplitude matched for each pH level). Zn²⁺-potentiated and control ATP-responses were normalized to the amplitude of responses to 100 μM ATP, which at pH 5.5–8.0 is a supramaximal stimulus.

6.5, $24.6 \pm 1.6 \mu\text{M}$; pH 5.5, inactive. A similar pH-dependency was observed where P2X₂ receptors were pre-incubated with Zn²⁺ for 5 min (Figure 3b). Under these conditions, the apparent EC₅₀ values for the Zn²⁺ potentiation of I_{ATP} were ($n=4$): pH 8.0, $7.4 \pm 1.1 \mu\text{M}$; pH 7.5, $6.1 \pm 1.2 \mu\text{M}$; pH 6.5, $35.3 \pm 4.7 \mu\text{M}$; pH 5.5, inactive.

Effect of Zn²⁺ on concentration-dependence of I_{ATP}

When applied simultaneously with ATP (0.1–300 μM), 10 μM and 30 μM Zn²⁺ displaced the ATP concentration-response (C/R) curve leftwards, without changing the maximum response or the slope of the curve, while 100 μM and 300 μM Zn²⁺ did not significantly displace the ATP C/R curve further to the left (Figure 4a). Thus, the potentiating effect of Zn²⁺ reached a maximum at approximately 30 μM . EC₅₀ values for ATP, at the following Zn²⁺ concentrations, were ($n=4$): 0 μM , $16.2 \pm 1.4 \mu\text{M}$; 10 μM , $6.5 \pm 0.8 \mu\text{M}$; 30 μM , $3.1 \pm 0.33 \mu\text{M}$; 100 μM , $2.4 \pm 0.1 \mu\text{M}$; 300 μM , $3.3 \pm 0.3 \mu\text{M}$.

Prolonged pre-incubation with Zn²⁺ had an inhibitory effect on ATP activity and also reduced ATP potency. Where P2X₂ receptors were pre-incubated with 100 μM Zn²⁺ for 60 min ($T=60$ min), the ATP C/R curve was displaced to the right of the control curve and maximum ATP activity was reduced by about 50% (Figure 4b). This inhibition was reversed on washout. However, without pre-incubation 100 μM Zn²⁺ applied simultaneously with ATP enhanced agonist potency and displaced the ATP-curve to the left of the control curve (Figure 4b). EC₅₀ values for ATP, at the following Zn²⁺ concentrations and incubation times, were ($n=4$): 0 μM , $16.2 \pm 1.4 \mu\text{M}$; 100 μM ($T=0$), $2.4 \pm 0.1 \mu\text{M}$; 100 μM ($T=60$), $50.9 \pm 8.0 \mu\text{M}$. The inhibitory effect of prolonged Zn²⁺ pre-incubation on ATP was prevented by lowering pH to 5.5 (Figure 4c). EC₅₀ values (at pH 5.5) for ATP, at the following Zn²⁺ concentrations and incubation times, were ($n=4$): 0 μM , $547 \pm 39 \text{ nM}$; 100 μM ($T=0$), $577 \pm 89 \text{ nM}$; 100 μM ($T=60$), $526 \pm 86 \text{ nM}$ (none significantly different).

Zn²⁺ and pH interactions on concentration-dependence of I_{ATP}

Acidic shifts have been shown to increase ATP potency at P2X₂ receptors (King *et al.*, 1996c; Wildman *et al.*, 1997), with an apparent pK_a of 7.05 ± 0.05 (King *et al.*, 1997). To determine any interaction between Zn²⁺ and pH, the concentration-dependence of I_{ATP} was investigated at pH 7.5, 6.5 and 5.5 in the absence and presence of 100 μM Zn²⁺ (Figure 5a,b,c). With H⁺ alone, ATP potency at P2X₂ receptors was increased at pH 6.5 and reached a maximum at pH 5.5. EC₅₀ values for ATP were ($n=4$): pH 7.5, $16.2 \pm 1.4 \mu\text{M}$; pH 6.5, $1.38 \pm 0.22 \mu\text{M}$; pH 5.5, $547 \pm 39 \text{ nM}$. Thus, ATP potency was increased 12 fold at pH 6.5 and 30 fold at pH 5.5. Where Zn²⁺ (100 μM) was applied simultaneously with ATP, increases in ATP potency were progressively reduced as pH was lowered to 5.5. Thus, EC₅₀ values for ATP (Zn²⁺ present) were ($n=4$): pH 7.5, $2.4 \pm 0.13 \mu\text{M}$; pH 6.5, $363 \pm 43 \text{ nM}$; pH 5.5, $481 \pm 79 \text{ nM}$. Thus, Zn²⁺ increased the ATP potency at P2X₂ receptors by 7 fold at pH 7.5 and 4 fold at pH 6.5, but had no further effect at pH 5.5.

Effect of DEPC on Zn²⁺ and H⁺ potentiation

The imidazole group on histidyl residues shows a strong avidity for both Zn²⁺ and H⁺ ions and is irreversibly denatured by diethylpyrocarbonate (DEPC) to abolish Zn²⁺




Figure 4 Effects of Zn²⁺ on concentration-dependence of I_{ATP} . In (a), concentration-dependence of whole-cell membrane currents to ATP (0.1–300 μM , at pH 7.5) in the presence of Zn²⁺ applied at the same time as the agonist. In (b) and (c), effect of Zn²⁺ (100 μM) applied at the same time as ATP ($T=0$) or continuously over a 60 min pre-incubation ($T=60$) on the concentration-dependence of ATP-responses at pH 7.5 (b) and pH 5.5 (c). Concentration-response curves to ATP in the absence of Zn²⁺ are also shown.

and H⁺ binding (Miles, 1977). DEPC was applied briefly (10 min) in single doses (0.3, 0.5, 1 or 3 mM) to *Xenopus* oocytes ($n = 3$, per dose) expressing the P2X₂ receptors to see if

the potentiating effects of Zn²⁺ and H⁺ depended on histidyl residues near the ATP binding site. At concentrations greater than 0.3 mM, DEPC reduced maximal ATP-responses by 90% and decreased ATP potency in an irreversible manner. In spite of this inhibitory action by DEPC, residual ATP-responses were potentiated by Zn²⁺ (100 μM) and pH (6.5). Histidyl residues did not appear to be crucial for the potentiating effect of Zn²⁺ and H⁺ on I_{ATP} .

Effect of Zn²⁺ on suramin blocking activity

The blocking activity of suramin at P2X₂ receptors is progressively increased by acidic shifts in extracellular pH (King et al., 1997). Zn²⁺ (100 μM) mimicked this action of H⁺ ions and enhanced blocking activity of suramin when Zn²⁺ was applied simultaneously with ATP (Figure 6). IC₅₀ values for suramin at the following Zn²⁺ concentrations were ($n = 4$): 0 μM, 1.74 ± 0.17 μM; 100 μM, 225 ± 23 nM.

Discussion

Extracellular Zn²⁺ and acidic solutions reversibly potentiate I_{ATP} at P2X₂ receptors, an effect caused by a leftwards displacement of the ATP concentration-response curve and increase in agonist potency. The potentiating action of Zn²⁺ and H⁺ was saturable, reaching a maximum at 30 μM and pH 5.5 (i.e. [H⁺] = 30 μM), respectively. Zn²⁺ and H⁺ also enhanced the blocking activity of suramin at P2X₂ receptors, thus showing that agonist and antagonist activity are both modulated and raising the possibility that Zn²⁺ and H⁺ might act through a common mechanism. It is unlikely that these potentiating effects were due to an inhibitory action on ecto-ATPases, since the modulatory actions of Zn²⁺ and H⁺ on ATP activity differ in time-course and the defolliculated oocyte




Figure 5 Interaction of Zn²⁺ and H⁺ on concentration-dependence of I_{ATP} . Concentration-dependence of whole-cell membrane currents to ATP (0.03–300 μM) at pH 7.5 (a), pH 6.5 (b) and pH 5.5 (c), in the absence and presence of Zn²⁺ (100 μM) applied at the same time as the agonist. The ATP C/R curve was maximally displaced to the left by 100 μM Zn²⁺ at pH 7.5 (see (a); also Figure 4a). The control ATP C/R curve at pH 7.5 is shown as the dashed curve in (b) and (c).




Figure 6 Zn²⁺ modulation of suramin antagonism of I_{ATP} . Concentration-dependence of suramin blockade of whole-cell membrane currents to ATP, in the absence and presence of Zn²⁺ (100 μM), applied at the same time as the agonist. Control ATP-responses were evoked with the EC₃₀ (i.e. control ATP-responses were amplitude matched to account for the potentiating effect of Zn²⁺) and the blocking activity of suramin tested. Experiments were carried out at pH 7.5.

is largely devoid of surface enzymes that degrade ATP (Ziganshin *et al.*, 1995). Furthermore, the potentiating effects of Zn²⁺ and H⁺ cannot be attributed to an interaction with ATP itself, since low pH levels and Zn²⁺ do not universally enhance ATP activity at all of the subtypes of cloned P2X receptors.

H⁺ exerted a greater effect than Zn²⁺ on ATP-responses at P2X₂ receptors. For acid shifts, ATP potency was maximally increased 30 fold with the EC₅₀ values of 547 ± 39 nM at pH 5.5 and 16.2 ± 1.4 μM at pH 7.5. However, for the transition metal ATP potency was maximally increased 7 fold with an EC₅₀ value of 2.4 ± 0.1 μM in the presence of 100 μM Zn²⁺ compared to a control value of 16.2 ± 1.4 μM (at pH 7.5). The potentiating effects of H⁺ persisted with time while Zn²⁺ acted in a time-dependent manner and potentiation showed run-down and, eventually, was replaced by an inhibitory action. The greater effectiveness and persistence of H⁺ potentiation may help explain why the inhibitory actions of prolonged Zn²⁺ pre-incubation were reversed when pH_e was lowered to 5.5, a physiological antagonism of Zn²⁺ inhibition by a stronger and opposing H⁺ potentiating effect. We also noticed that ATP potency decreases when the pH of the superfusate is raised from 7.5 to 8.0, while the extent of Zn²⁺ potentiation was no different at pH 7.5 and pH 8.0. This finding, together with the above differences in effectiveness of Zn²⁺ and H⁺, raised a question mark over these modulators acting at the same allosteric site on the P2X₂ receptor.

Others have briefly commented on the potentiating effect of Zn²⁺ (Brake *et al.*, 1994; Nakazawa & Ohno, 1996; 1997). Brake *et al.* (1994) tested 10 μM Zn²⁺ against I_{ATP} at P2X₂ receptors and found that this group IIb transition metal increased agonist potency by 4 fold. Here, we found a 3 fold increase in agonist potency under similar conditions. Nakazawa & Ohno (1996, 1997) also found that 100 μM Zn²⁺ was less effective than 10 μM at potentiating ATP-responses at P2X₂ receptors, confirming that the Zn²⁺ C/R curve is bell-shaped. However, this appears to be true only when P2X₂ receptors are pre-incubated with this transition metal.

The present results on H⁺ potentiation of I_{ATP} agree with earlier findings on pH sensitivity of P2X₂ receptors (King *et al.*, 1996c; 1997; Wildman *et al.*, 1997), except for 2 minor differences. Previously, we found no significant difference in EC₅₀ values for ATP at pH 6.5 and pH 5.5 (1.2 ± 0.1 μM and 1.1 ± 0.1 μM, respectively) (King *et al.*, 1996c; 1997). However, a small difference in EC₅₀ values was shown in the present study, i.e. 1.38 ± 0.22 μM (at pH 6.5) and 547 ± 39 nM (at pH 5.5) (*P* < 0.05). We explain this difference solely on a greater accuracy in determining EC₅₀ values. We also found that an original estimate of suramin activity (IC₅₀ = 10.4 ± 1.2 μM)

(King *et al.*, 1997) did not agree with the present data (IC₅₀ = 1.74 ± 0.17 μM). The blocking activity of suramin is exceedingly sensitive to pH_e (King *et al.*, 1997) and, therefore, small differences in setting pH_e have a profound impact on determinations of IC₅₀ values. Also, we are now aware that the life-time and accuracy of pH-probes are often less than stated by their manufacturers.

Histidyl residues have been implicated in the effects of Zn²⁺ and H⁺ on agonist activity/binding at adenosine and γ-aminobutyric acid (GABA) receptors (Smart, 1990; Allende *et al.*, 1993; Wang *et al.*, 1995; Krishek 1996). We used DEPC to denature histidyl residues on the extracellular loop of the P2X₂ receptor, although this reagent is not selective for histidine and will also act on arginyl, sulphhydryl and tyrosyl residues when used at high concentrations (10 mM) and under very acidic conditions (pH 4) (Miles, 1977). DEPC did not block the potentiating effects of Zn²⁺ and H⁺ at P2X₂ receptors and this outcome appeared to discount the involvement of histidyl residues as the allosteric sites near the ATP binding site. The precise locus of these sites on P2X₂ remains to be determined.

Cloues (1996) showed that the native P2X receptors on SCG neurones, PC12 cells and sensory neurones in rat all have similar biophysical and pharmacological properties. The P2X receptors at these three sites are weakly stimulated by ADP (Rhoads *et al.*, 1993; Khakh *et al.*, 1995) and, interestingly, the P2X₂ receptor is also insensitive to ADP (King *et al.*, 1997). ATP-responses at the above native P2X receptors are strongly affected by Zn²⁺ (Li *et al.*, 1993; Koizumi *et al.*, 1995) as are ATP-responses at the P2X₂ receptor. Acidic shifts potentiate and alkaline shifts inhibit ATP-responses at native P2X receptors of sensory neurones (Li *et al.*, 1996) and, to date, only the P2X₂ receptors have been shown to react in the same way to pH changes (King *et al.*, 1997; North, unpublished data). Thus, it appears that weak ADP activity, Zn²⁺ potentiation of ATP-responses and sensitivity to pH_e may represent key pharmacological features that could identify the presence of P2X₂ subunits in native P2X receptors. It is significant, therefore, that P2X₂ transcripts are contained in PC12 cells, sympathetic (SCG and coeliac) neurones and sensory (nodose and DRG) neurones (Brake *et al.*, 1994; Collo *et al.*, 1996).

This work was supported by the British Heart Foundation and Institut de Recherches Servier (France). We are grateful to Dr David Julius (UCSF, U.S.A.) for the gift of cDNA encoding the rat P2X₂ receptor. We thank Dr Alan North (Glaxo, Geneva) for information on pH sensitivity of P2X_{1,4} receptors.

References

- ALLENDE, G., CASADÓ V., MALLOL, J., FRANCO, R., LLUIS, C. & CANELA, E. (1993). Role of histidine residues in agonist and antagonist binding sites of A₁ adenosine receptor. *J. Neurochem.*, **60**, 1525–1533.
- BRAKE, A.J., WAGENBACH, M.J. & JULIUS, D. (1994). New structural motif for ligand gated ion channels defined by an ionotropic ATP receptor. *Nature*, **371**, 519–523.
- BUELL, G., COLLO, G. & RASSENDREN, F. (1996). P2X receptors: an emerging channel family. *Eur. J. Neurosci.*, **8**, 2221–2228.
- BURNSTOCK, G. (1996). P2 purinoceptors: historical perspective and classification. In: *P2 purinoceptors: Localization, Function and Transduction Mechanisms*, pp. 1–34, Chichester (Ciba Foundation Symposium, 198); John Wiley & Sons.
- BURNSTOCK, G. & KING, B.F. (1996). Numbering of cloned P2 purinoceptors. *Drug Devel. Res.*, **38**, 67–71.
- CLOUES, R. (1995). Properties of ATP-gated channels recorded from rat sympathetic neurons; voltage dependence and regulation by Zn²⁺ ions. *J. Neurophysiol.*, **73**, 312–319.
- CLOUES, R., JONES, S. & BROWN, D.A. (1993). Zn²⁺ potentiates ATP-activated currents in rat sympathetic neurons. *Pflügers Arch.*, **424**, 152–158.
- COLLO, G., NORTH, R.A., KAWASHIMA, E., MERLO-PICH, E., NEIDHART, S., SURPRENANT, A. & BUELL, G. (1996). Cloning of P2X₅ and P2X₆ receptors and the distribution and properties of an extended family of ATP-gated ion channels. *J. Neurosci.*, **16**, 2495–2507.
- KHAKH, B.S., HUMPHREY, P.P.A. & SURPRENANT, A. (1995). Electro-physiological properties of P2X-purinoceptors in rat superior cervical ganglion, nodose and guinea-pig coeliac neurones. *J. Physiol.*, **484**, 385–395.

- KING, B.F., PINTOR, J., WANG, S., ZIGANSHIN, A.U., ZIGANSHINA, L.E. & BURNSTOCK, G. (1996a). A novel P1 purinoceptor activates an outward K⁺ current in follicular oocytes of *Xenopus laevis*. *J. Pharmacol. Exp. Ther.*, **276**, 93–100.
- KING, B.F., WANG, S. & BURNSTOCK, G. (1996b). P2 purinoceptor-activated inward currents in follicular oocytes of *Xenopus laevis*. *J. Physiol.*, **494**, 17–28.
- KING, B.F., WILDMAN, S.S., ZIGANSHINA, L.E., PINTOR, J. & BURNSTOCK, G. (1997). Effects of extracellular pH on agonism and antagonism at a recombinant P2X₂ receptor. *Br. J. Pharmacol.*, **121**, 1445–1453.
- KING, B.F., ZIGANSHINA, L.E., PINTOR, J. & BURNSTOCK, G. (1996c). Full sensitivity of P2X₂ purinoceptor to ATP revealed by changing extracellular pH. *Br. J. Pharmacol.*, **117**, 1371–1373.
- KOIZUMI, S., IKEDA, M., INOUE, K., NAKAZAWA, K. & INOUE, K. (1995). Enhancement by zinc of ATP-evoked dopamine release from rat pheochromocytoma PC12 cells. *Brain Res.*, **673**, 75–82.
- KRISHEK, B.J., AMATO, A., CONNOLLY, C.N., MOSS, S.J. & SMART, T.G. (1996). Proton sensitivity of the GABA_A receptor is associated with the receptor subunit composition. *J. Physiol.*, **492**, 431–443.
- LEWIS, C., NEIDHART, S., HOLY, C., NORTH, R.A., BUELL, G. & SURPRENANT, A. (1995). Co-expression of P2X receptor subunits can account for ATP-gated currents in sensory neurons. *Nature*, **377**, 432–435.
- LI, C., PEOPLES, R.W., LI, Z. & WEIGHT, F.F. (1993). Zn²⁺ potentiates excitatory action of ATP on mammalian neurons. *Proc. Natl. Acad. Sci. U.S.A.*, **90**, 8264–8267.
- LI, C., PEOPLES, R.W. & WEIGHT, F.F. (1996). Proton potentiation of ATP-gated ion channel responses to ATP and Zn²⁺ in rat nodose ganglion neurons. *J. Neurophysiol.*, **76**, 3048–3058.
- MILES, E.W. (1977). Modification of histidyl residues in proteins with diethylpyrocarbonate. *Methods Enzymol.*, **47**, 431–442.
- NAKAZAWA, K. & OHNO, Y. (1996). Dopamine and 5-hydroxytryptamine selectively potentiate neuronal type ATP receptor channels. *Eur. J. Pharmacol.*, **296**, 119–122.
- NAKAZAWA, K. & OHNO, Y. (1997). Effects of neuroamines and divalent cations on cloned and mutated ATP-gated channels. *Eur. J. Pharmacol.*, **325**, 101–108.
- NORTH, R.A. (1996). Families of ion channels with two hydrophobic segments. *Curr. Opin. Cell Biol.*, **8**, 474–483.
- RHOADS, A.R., PARUI, R., VU, N.D., CADOGAN, R. & WAGNER, P.D. (1993). ATP-induced secretion in PC12 cells and photoaffinity labeling of receptors. *J. Neurochem.*, **61**, 1657–1666.
- SMART, T.G. (1990). Uncultured lobster muscle, cultured neurons and brain slices: the neurophysiology of zinc. *J. Pharm. Pharmacol.*, **42**, 377–387.
- WANG, T.L., HACKAM, A., GUGGINO, W.B. & CUTTING, G.R. (1995). A single histidine residue is essential for zinc inhibition of GABA ρ1 receptors. *J. Neurosci.*, **15**, 7684–7691.
- WILDMAN, S.S., KING, B.F. & BURNSTOCK, G. (1997). Potentiation of ATP-responses at a recombinant P2X₂ receptor by neurotransmitters and related substances. *Br. J. Pharmacol.*, **120**, 221–224.
- ZIGANSHIN, A.U., ZIGANSHINA, L.E., KING, B.F. & BURNSTOCK, G. (1995). Characteristics of ecto-ATPase of *Xenopus* oocyte and the inhibitory actions of suramin on ATP breakdown. *Pflügers Arch.*, **429**, 412–418.

(Received September 17, 1997

Revised December 3, 1997

Accepted December 8, 1997)